

# Pharmaceutical Care Services and Results in Project ImPACT: Hyperlipidemia

Benjamin M. Bluml, James M. McKenney, and Mark J. Cziraky

**Objective:** To demonstrate that pharmacists, working collaboratively with patients and physicians and having immediate access to objective point-of-care patient data, promote patient persistence and compliance with prescribed dyslipidemic therapy that enables patients to achieve their National Cholesterol Education Program (NCEP) goals. **Design:** Observational study. **Participants:** 26 community-based ambulatory care pharmacies: independent, chain-professional, chain-grocery store, home health/home infusion, clinic, health maintenance organization/managed care. **Main Outcome Measures:** Rates of patient persistence and compliance with medication therapy and achievement of target therapeutic goals. **Results:** In a population of 397 patients over an average period of 24.6 months, observed rates for persistence and compliance with medication therapy were 93.6% and 90.1%, respectively, and 62.5% of patients had reached and were maintained at their NCEP lipid goal at the end of the project. **Conclusion:** Working collaboratively with patients, physicians, and other health care providers, pharmacists who have ready access to objective clinical data, and who have the necessary knowledge, skills, and resources, can provide an advanced level of care that results in successful management of dyslipidemia.

*J Am Pharm Assoc. 2000;40:157-65.*

Project ImPACT: Hyperlipidemia, a community pharmacy-based demonstration project, was initiated in March 1996 and completed in October 1999. *ImPACT* is an acronym for Improve Persistence And Compliance with Therapy.

Dyslipidemia (hyperlipidemia) was considered an ideal area in which to demonstrate the value that pharmacists can add to the patient care process for several reasons:

- Coronary artery disease (CAD) is the leading cause of death in the United States and accounts for an annual expenditure of \$100 billion for health care.<sup>1</sup>
- Dyslipidemia has been shown to be associated with increased risk of CAD in large epidemiologic studies.<sup>2</sup>
- Reduction in low-density lipoprotein cholesterol (LDL-C) levels has been shown to produce reductions in CAD events and total mortality.<sup>3-9</sup>
- Other modifiable CAD risk factors are invariably present in

patients with hyperlipidemias, including hypertension, diabetes, obesity, and sedentary lifestyle.

- Pharmacist services are widely accessible to patients, physicians, and other health care providers and add a unique pharmacotherapy management resource to the health care delivery team.
- Evidence suggests that pharmacists who provide disease management services can increase patient compliance and improve treatment outcomes.<sup>10-12</sup>
- A point-of-care testing device for measuring lipid levels, the Cholestech LDX Analyzer, is available to pharmacists and other health care providers.
- The availability of reliable patient lipid profile results within 5 minutes of obtaining a blood sample by fingerstick allows the pharmacist to be directly involved in management of lipid-lowering therapies and patient outcomes.
- The management of cholesterol disorders represents a major benchmark by which quality health care services can be evaluated by accrediting agencies and purchasers of health care.<sup>13</sup>

Lifestyle modifications combined with improvements in persistence and compliance in the use of lipid-lowering medications will result in a greater number of patients reaching their target lipid goals. If patients reach and maintain their National Cholesterol Education Program (NCEP) goals, cardiovascular-related risk will be reduced,<sup>6,7</sup> resulting in positive health care outcomes. Table 1 provides an overview of desirable ranges for selected lipid measures as recommended by the NCEP.<sup>3</sup>

Received December 22, 1999, and in revised form January 20, 2000. Accepted for publication February 1, 2000.

Benjamin M. Bluml is vice president for research, APhA Foundation, Washington, D.C. James M. McKenney, PharmD, is professor, School of Pharmacy, Virginia Commonwealth University, Richmond. Mark J. Cziraky, PharmD, is executive vice president and chief operating officer, Health Core, Inc., Newark, Del.

Correspondence: Benjamin M. Bluml, 2215 Constitution Avenue, NW, Washington, DC 20037-2985. Fax: 202-429-6300. E-mail: bmb@mail.aphanet.org.

See related Viewpoint on page 143.

**Table 1. Desirable Ranges for Selected Lipid Measures**

Total Cholesterol	Triglycerides	HDL-C	LDL-C <sup>a</sup>
< 200 mg/dL desirable	< 200 mg/dL desirable	≥ 35 mg/dL desirable	< 160 mg/dL goal if < 2 risk factors
200–239 mg/dL borderline	200–400 mg/dL borderline	< 35 mg/dL low	< 130 mg/dL goal if ≥ 2 risk factors
≥ 240 mg/dL high	400–1,000 mg/dL high		≤ 100 mg/dL goal if CAD history
	> 1,000 mg/dL very high		

CAD = coronary artery disease; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol.

<sup>a</sup>National Cholesterol Education Program Guidelines.

## Objectives

The core objectives of Project ImPACT: Hyperlipidemia were to:

1. Improve patient persistence and compliance with lipid-lowering therapy.
2. Increase communication and the flow of clinical information among patients, pharmacists, and physicians.
3. Improve the cholesterol levels of individual patients over time.
4. Increase the population of patients who reach and maintain their NCEP lipid goals.

## Methods

### Site Selection

As the result of a competitive application process, 32 community pharmacy practice sites distributed across 15 states were selected to participate in Project ImPACT: Hyperlipidemia. Selection was based on criteria that addressed the readiness of the pharmacy to provide basic pharmaceutical care services as evidenced by the availability of certain health care resources and the requisite knowledge and skills to facilitate the delivery of such services:

- Private or semiprivate area for patient consultation.
- Technician support.
- Documentation system for recording, tracking, and reporting patient care interventions.
- Experience with patient-focused disease state management programs.
- Demonstrated communication skills.
- Ability to implement point-of-care testing technologies.

In addition, participating pharmacists from all sites attended a 2½-day orientation and training program at the project's inception. That training program was the basis for the APhA certificate program "Pharmaceutical Care for Patients with Dyslipidemias."

Of the 32 pharmacies, 2 sites were unable to implement the project (one secondary to regulatory issues, and the other secondary to departmental reorganization), 2 experienced unexpected staffing challenges, 1 moved to another location, and 1 pharmacy was sold and closed. Thus, a total of 26 pharmacy practice sites in 12 states completed the study (Table 2).

### Patient Enrollment

Patients enrolled in the project were either newly diagnosed with dyslipidemia (e.g., hypercholesterolemia, mixed hyperlipidemia) or were already receiving lipid-lowering medications but were poorly controlled (i.e., not yet at target lipid goal). Patients were identified through referrals from local physicians or other health care providers, by the project pharmacists, or by patient self-referral. In cases of nonphysician referral, patients' physicians were contacted by pharmacists and were involved from that point forward in the patient's care. Patients were informed about the expected effects of their participation in the project (i.e., potential benefits, risks, inconveniences, discomforts), were assured confidentiality (patient privacy was protected by use of an assigned code in all reporting), and told about their right to withdraw at any time. Patients gave written informed consent and authorized that medical information from other health care providers could be sent to the pharmacist.

### Process of Care

Patients provided the necessary personal and general health information that the pharmacist used to assess their CAD risk. From a fingerstick blood sample, a fasting lipid profile was obtained using the Cholestech LDX Analyzer (a point-of-care testing device in the "waived" category under the Clinical Laboratory Improvement Amendments<sup>14</sup>), and results were logged into a clinical activity record at each project visit. After the initial visit and consultation with the pharmacist, patients were asked to make follow-up visits every month for the first 3 months and quarterly thereafter. In addition to being actively involved in their therapy, treatment plans, and goal setting, patients as well as their physicians were kept informed about clinical progress in these areas:

- Cholesterol test results.
- Condition.
- CAD risk.
- NCEP goal achievement.

### Practice Model

The practice model designed for the project was sufficiently flexible to accommodate variations in staffing and types of resources available at the practice sites represented in the study.

**Table 2. Project ImPACT: Hyperlipidemia Participating Pharmacies by Practice Setting**

Community Pharmacy Practice Setting	No.
<b>Independent</b>	<b>14</b>
Baggett Pharmacy (Levelland, Tex.)	
Bel-Aire Pharmacy (White Bear Lake, Minn.)	
Goodrich Pharmacy (Anoka, Minn.)	
Lutz Pharmacy (Altoona, Iowa)	
Mar-Main Pharmacy (South Bend, Ind.)	
Mullins Drugs (Birmingham, Ala.)	
Northaven Pharmacy (Seattle, Wash.)	
Osterhaus Pharmacy (Maquoketa, Iowa)	
QFC Pharmacy (Seattle, Wash.)	
Red Wing Corner Drug (Red Wing, Minn.)	
Ritzman Pharmacy (Akron, Ohio)	
Travis Pharmacy, Inc. (Shenandoah, Iowa)	
Uptown Pharmacy (Westerville, Ohio)	
West End Drug (Bar Harbor, Maine)	
<b>Chain—professional</b>	<b>3</b>
Medicap Pharmacy (Urbandale, Iowa)	
Medicap Pharmacy (Wilmington, N.C.)	
Medicine Shoppe Pharmacy (Youngstown, Ohio)	
<b>Chain—grocery store</b>	<b>1</b>
Ukrop's Pharmacy (Richmond, Va.)	
<b>Home health/home infusion</b>	<b>2</b>
Bohlman Drug Store, Inc. (Boscobel, Wisc.)	
Jones Pharmacy & Home Health Care (Spokane, Wash.)	
<b>Clinic pharmacy</b>	<b>4</b>
Family PharmaCare Center, Inc. (West Lafayette, Ind.)	
Goodrich Pharmacy (Elk River, Minn.)	
Hadfield's Pharmacy (Edmonds, Wash.)	
Pharmaceutical Care Clinic, Ohio State University (Columbus, Ohio)	
<b>Health maintenance organization/managed care</b>	<b>2</b>
Health Core, Inc. (Newark, Del.)	
Kaiser Foundation Health Plan Pharmacy (Cleveland, Ohio)	

ImPACT = Improve Persistence And Compliance with Therapy.

The practice model structure:

- Established a process for the seamless flow of patient care data between and among patients, pharmacists, and physicians.
- Used point-of-care testing technology to obtain timely, objective information about the patient's progress in a community practice setting.
- Organized methods for pharmacists to document, interpret, and report their lipid management interventions.

Figure 1 depicts the collaborative care process that provided the framework for the Project ImPACT practice model.

## Persistence and Compliance Measures

The persistence measure used for the project was defined as follows: a patient who started on medication, remained on medication subsequent to drug therapy initiation, and continued to be on medication as of his or her last visit. Persistence as a percentage was calculated by dividing the number of persistent patients by the total number of patients who started on medication. Com-

pliance was determined through an evaluation based on the number of missed doses for each lipid-lowering medication and refill timing. Any patient who missed doses for 5 days or more or who missed a scheduled refill visit by more than 5 days was deemed to be noncompliant at that visit. Compliance as a percentage was calculated by dividing the number of visits at which patients were compliant by the total number of patient visits.

## End-of-Project Survey

A final project survey was conducted with all 26 sites to gain an understanding of what factors were likely responsible for creating the environment that produced the persistence, compliance, and treatment-to-goal results. The survey also included queries about the sites' experiences with obtaining payment for the pharmaceutical care services delivered during this project.

## Results

A total of 574 patients were enrolled at the 26 sites before July 1, 1997. Of those, 397 patients completed the entire study, and results are presented for those patients in the following section. There were 34 patients who completed only 1 visit and had insufficient data to allow reporting of results. The remaining 143 patients did complete at least 2 visits to the pharmacy, but did not complete the full 2-year observation period: 29 withdrew in the first 90 days, 30 moved from the area, 33 gave personal reasons, 22 had logistical or medical complications, and 29 were lost to follow-up. The results for these 143 patients are reported separately at the end of the Results section.


## Patient Population Characteristics

At the beginning of the study, 153 (38.5%) of the 397 patients were either newly diagnosed or had been taking lipid-lowering medications for less than 1 month, and 244 (61.5%) had been on lipid-lowering medications for longer periods but remained poorly controlled. This combined population consisted of 51.6% women and 48.4% men, with an average age of 57 years. Of these patients, 298 (75.1%) had no history of CAD and were categorized as primary prevention patients (199 with an LDL-C goal < 130 mg/dL and 99 with an LDL-C goal < 160 mg/dL), while the other 99 (24.9%) had previously experienced a coronary event and therefore fell into a secondary prevention category (LDL-C goal ≤ 100 mg/dL). Patient ethnicity was as follows: 24 (6%) African American, 3 (0.8%) Asian, 337 (84.9%) Caucasian, 1 (0.2%) Hispanic, and 32 (8.1%) not specified.

## Persistence and Compliance with Medication Therapy

Of the 397 patients who completed the 2-year study,

Figure 1. Project IMPACT Collaborative Care Process



CAD = coronary artery disease; IMPACT = Improve Persistence And Compliance with Therapy.

345 (86.9%) patients were treated with lipid-lowering medications and lifestyle modifications, while 52 (13.1%) continued with lifestyle modifications focused on diet and exercise in an effort to reach target cholesterol goals. The distribution of lipid-lowering medication use was as follows:

- 89% HMG-CoA reductase inhibitors.
- 5% niacin.
- 4% fibrates.
- 2% bile acid resins.

Of the 345 patients started on medication, 323 continued with drug therapy, for a resultant patient medication persistence rate of 93.6%.

Of 2,817 documented visits for patients on medications, 2,539 occurrences of compliance (i.e., within 5 days of expected refills) were reported, for a resulting per-visit medication compliance rate of 90.1%.

### Resultant Lipid Levels

Using the two-tailed Student *t* test for paired data, statistically significant improvements were found for the 397 study patients using beginning and ending LDL-C measures (mean duration = 24.6 months; Table 3). Mean ( $\pm$  standard deviation) reductions of  $12.8\% \pm 1.6\%$  and  $10.0\% \pm 6.5\%$  were observed for total cholesterol and triglycerides, respectively, while a mean increase of  $14.2\% \pm 3.9\%$  was observed in high-density lipoprotein levels. Overall, a mean reduction of  $22.1\% \pm 2.6\%$  was observed for LDL-C values. In addition, the midpoint measures (mean interval from beginning = 12.1 months; Table 3) demonstrate progressive improvements over time.

The NCEP Adult Treatment Panel II (ATPII) guidelines recommend LDL-C goals of  $< 160$  mg/dL for patients with less than two CAD risk factors,  $< 130$  mg/dL for patients with two or more CAD risk factors, and  $\leq 100$  mg/dL for patients with a history of CAD (see Table 1). Based on these NCEP guidelines, 290 of the 397 patients (73.1%) were at or below goal on two or more visits during the study, and 248 (62.5%) were at or below goal as of their last full lipid profile. Figure 2 depicts NCEP goal achievement at the end of the project in the primary and secondary prevention groups and in the combined patient population.

### Pharmacist Interventions

Pharmacists intervened with physicians to request a variety of therapeutic changes during the course of the study. These interventions were focused on improving NCEP goal achievement through drug therapy optimization and addressed issues that included coordination of care, adverse drug reactions, drug interactions, drug dosing, drug selection, and side effects. Physicians accepted the pharmacist recommendations and made changes in 265 (76.6%) of the 346 reported interventions.

### Practice Model Observations

While implementation of the Project ImPACT practice model may have varied slightly from site to site to accommodate practice differences, it consistently produced an environment that resulted in a high level of collaboration through the following:

- Regular communications between and among all involved parties.
- Referral of patients by pharmacists to physicians and other providers (family practitioners, internal medicine physicians, cardiologists, dietitians, nurse practitioners, and endocrinologists).
- Referral of patients to pharmacists by physicians and other providers (family practitioners, internal medicine physicians, cardiologists, and nurse practitioners).
- Increased availability and use of objective clinical measures.
- Sharing treatment data and pertinent lifestyle and clinical information, including objective lipid measures obtained in the pharmacy, with patients and physicians.
- Periodic evaluation of the patient's progress toward lipid goals, and, if necessary, consultation and intervention with the patient's physician.
- Timely adjustments in the patient's treatment plans.

### Process of Care Observations

Eligible "at risk" patients who were enrolled in the project were identified through community screening events (12%), patient self-referrals (13%), physician referrals (15%), and pharmacist identification and referral (60%).

Two critical components of the process of care in the pharmacy were scheduling appointments for patients and arranging for ade-


**Table 3. Fasting Mean Lipid Levels of Patients in Study**

Measure (No. Patients)	Beginning Measure mg/dL (SD)	Midpoint Measure mg/dL (SD)	Ending Measure mg/dL (SD)	Mean Change	<i>P</i> value <sup>a</sup>
Total cholesterol (396)	238.0 (46.7)	216.7 (44.3)	207.5 (41.1)	-30.5 (3.9)	< .0001
Triglycerides (394)	216.6 (111.3)	199.1 (95.7)	195.0 (91.3)	-21.6 (14.1)	< .0001
High-density lipoproteins (394)	43.1 (14.1)	46.6 (15.6)	49.2 (16.5)	+6.1 (1.7)	< .0001
Low-density lipoproteins (387)	153.7 (41.3)	130.4 (37.8)	119.8 (35.7)	-33.9 (4.0)	< .0001

SD = standard deviation.

<sup>a</sup>Change calculated as ending measure less the beginning measure (mean duration = 24.6 months) and compared using a two-tailed Student *t* test for paired data.

**Figure 2. Percentage of Patients Who Achieved Their NCEP Goals During Study**



IMPACT = Improve Persistence And Compliance with Therapy; LDL-C = low-density lipoprotein cholesterol; NCEP = National Cholesterol Education Program.

quate personnel to provide the services. The end-of-project survey asked about the mechanisms that pharmacists used to accomplish these tasks. Table 4 lists the scheduling mechanisms and staffing arrangements used to accommodate the increased time commitment needed for the project. The right column indicates the percentage of sites in which those accommodations were made.

Because of the time management challenges that pharmaceutical care services can and often do present, the survey asked about the amount of time spent for the initial visit and for scheduled follow-up visits. On average, pharmacists spent 30 to 60 minutes (mean = 45 minutes) with patients at their initial visit and 10 to 30 minutes (mean = 22 minutes) with patients during follow-up visits.

Table 5 lists the various services that project sites used in managing the care of patients with dyslipidemia and the frequency with which these services or techniques were employed.

Pharmacists were asked to describe their level of satisfaction with their own role, their relationships with patients and physicians, and their perceptions of how satisfied patients and physicians were with pharmacists' services provided as a part of the project. The percentages of pharmacists responding "very satisfied" and "satisfied" were as follows:

- With their professional role, 88.5% and 11.5%, respectively.
- With their relationship with patients, 84.6% and 15.4%, respectively.
- With their relationship with physicians, 19.2% and 46.2%, respectively (with another 30.8% being "neutral" and 3.8% "dissatisfied").

Pharmacists perceived that 53.8% of their patients were "very satisfied" and 46.2% "satisfied" with the services provided. Pharmacists' perceptions of the physicians' feelings about the value of their services were not as strong: "very satisfied," 19.2%; "satisfied," 46.2%; "neutral," 26.9%; and "dissatisfied," 7.7%.

Pharmacists at 25 of the 26 project sites planned to continue to provide this service. Respondents at all sites recommended that

**Table 4. Scheduling and Staffing Arrangements in Study Pharmacies**

Description	% of Sites
Appointment scheduling	100
Before regular hours	80
Morning appointments	100
Afternoon appointments	88
Evening appointments	43
After regular hours	21
Weekend appointments	80
Added pharmacist staff	29
Increased pharmacist overlap	48
Changed pharmacist duties	72
Added technician staff	17
Increased technician overlap	5
Changed technician duties	59

other pharmacists implement these same types of services in their practices.

### Payment Observations

Although the project was not designed as a payment demonstration, participants were asked about the value of their services and their experiences in obtaining payment. Pharmacists indicated an average assigned value of \$55 per visit—\$28 for counseling services and \$27 for lipid profiles. With respect to patients paying for these services, pharmacists indicated that, of 232 patients who were asked for payment, 174 (75%) paid an average of \$35 per visit. Of 121 third party payers billed for services, 64 (53%) paid an average of \$30 for each visit billed. Of these 64 payers, 30 paid for counseling services and 53 paid for lipid profiles (some paid for both). Two project sites secured contracts with managed care organizations to deliver services to those health plan beneficiaries, one under a fee-for-service arrangement and the other under capitation.

### Patients Not Completing Study

The results presented thus far are for the population of patients who continued for the full duration of the project (Group 1). Data for those patients who did not complete the entire project (Group 2) show the following (Table 6):

- Patient demographics (age, ethnicity, sex, and CAD status) for Group 2 did not vary by more than 3% from Group 1.
- Average length of participation in the project was 7.2 months for Group 2, compared with 24.6 months for Group 1.
- Enrollment category distribution, newly diagnosed and poorly controlled, were 47.6% and 52.4%, respectively, in Group 2, and 38.5% and 61.5%, respectively, in Group 1.
- There were 20% fewer patients on drug therapy treatment in Group 2 as compared with Group 1.

- Persistence for Group 2 was 96.8%, compared with 93.6% for Group 1.
- Compliance for Group 2 was 86.1%, compared with 90.1% for Group 1.
- Clinical outcomes for lipid level and NCEP goal achievement measures for Group 2 were approximately 50% of those achieved by Group 1.

## Discussion

When evaluating the current state of dyslipidemia management in the existing health care delivery system, a less-than-optimal picture develops. Recent studies on the treatment of CAD indicate that the majority of eligible patients go untreated.<sup>15</sup> Of those patients who are treated, only 40% remain on their lipid-lowering medication therapy after 12 months.<sup>16-18</sup> Literature from primary care settings indicate that successful treatment-to-goal results range from 8% to 33%.<sup>19-21</sup>

The outcomes from Project ImPACT: Hyperlipidemia present a dramatically different picture. In the project, pharmacists demonstrated that they can, in collaboration with patients and physicians, effectively identify patients with lipid disorders who

require treatment and support them in their efforts to improve persistence, compliance, and treatment to goal. The results presented herein (see Figure 3), if compared with the existing health care delivery system, represent a twofold to fourfold improvement.

Project ImPACT: Hyperlipidemia provides a contemporary view of the capabilities of pharmacists, with the appropriate resources, to empower patients to achieve therapeutic outcomes through the effective application of a process of care to manage dyslipidemia. Pharmacists are in a prime position to ensure the success of collaborative practice efforts because of their accessibility to patients and physicians, their ability to use resources in providing an advanced level of care, and their information management capabilities, motivation to expand care, and education and training in the area of patient-focused disease management services. New point-of-care testing and communication technologies provide pharmacists with accurate, objective data to reinforce their counseling and intervention activities relative to persistence and compliance with diet, exercise, and drug therapy.

The project results suggest that patients receiving pharmaceutical care in a collaborative practice environment can make significant short-term improvements in persistence and compliance. However, longer-term participation in such an environment is required to achieve greater improvements in clinical outcomes.

**Table 5. Frequency of Use of Various Patient-Education Techniques in Study Pharmacies**

Service Provided	All or Most	Some	Rarely or Never
Explanation of the rationale for therapy	×		
Description of the dangers of atherosclerosis	×		
Education about the benefit of therapy in terms of reduced risk and enhanced chance of survival	×		
Explanation of how to interpret the lipid profile	×		
Identification of an LDL-C goal for the patient	×		
Identification of other health-related goals for the patient (e.g., weight, fat consumption)	×		
Showing patients a chart of their LDL-C results to monitor progress toward goal	×		
Praising patients for making progress toward (or achieving) their LDL-C (or other) goal	×		
Reinforcing the importance of compliance in reducing the risk of a heart attack	×		
Reviewing patients' compliance with therapy through such mechanisms as pill counts and refill records	×		
Questioning patients when noncompliance is detected or suspected	×		
Taking compliance histories during follow-up visits	×		
Answering questions patients have about atherosclerosis and its treatment	×		
Discussing the benefits and risks of lipid-lowering medications	×		
Encouraging patients to keep a log of the doses they take		×	
Encouraging patients to keep a log of their cholesterol results		×	
Helping patients solve problems in overcoming barriers to compliance		×	
Teaching a friend or family member about treatment in order to help them help the patient			×
Giving a tangible reward (e.g., coupon) for good compliance or achievement of goal			×
Giving patients ways to remind them of their doses (e.g., setting alarms, putting reminders on the refrigerator, putting the prescription bottle on the kitchen table)			×
Packaging the medication in ways to help patients take their drugs (e.g., organizers)			×
Calling patients at home to remind them of medication refills			×

LDL-C = low-density lipoprotein cholesterol.

**Table 6. Comparative Profile of Patient Data<sup>a</sup>**

Characteristic/Measure	Group 1 <sup>b</sup>	Group 2 <sup>c</sup>	All Patients
Total number of patients	397	143	540
Average age (years)	57.0	54.7	56.4
Ethnicity			
African American	24 (6)	8 (5.6)	32 (5.9)
Asian	3 (0.8)	0 (0)	3 (0.5)
Caucasian	337 (84.9)	123 (86)	460 (85.2)
Hispanic	1 (0.2)	1 (0.7)	2 (0.4)
Not specified	32 (8.1)	11 (7.7)	43 (8.0)
Sex			
Women	205 (51.6)	78 (54.5)	283 (52.4)
Men	192 (48.4)	65 (45.5)	257 (47.6)
CAD status			
Primary prevention	298 (75.1)	111 (77.6)	409 (75.7)
Secondary prevention	99 (24.9)	32 (22.4)	131 (24.3)
Project participation			
Average duration (months)	24.6	7.2	20
Enrollment category			
Newly diagnosed	153 (38.5)	68 (47.6)	221 (40.9)
Poorly controlled	244 (61.5)	75 (52.4)	319 (59.1)
Treatment			
Drug therapy and lifestyle modifications	345 (86.9)	95 (66.4)	440 (81.5)
Persistence with drug therapy (% patients)	93.6	96.8	94.3
Compliance with drug therapy (% patients)	90.1	86.1	89.7
Lifestyle modifications only	52 (13.1)	48 (33.6)	100 (18.5)
Clinical outcomes			
Total cholesterol (% change from baseline)	-12.8	-5.4	-10.8
Triglycerides (% change from baseline)	-10.0	-5.7	-8.7
HDL-C (% change from baseline)	+14.2	+6.5	+12.2
LDL-C (% change from baseline)	-22.1	-13.1	-19.8
NCEP goal achievement (% at ending measure)	62.5	38.5	56.1


CAD = coronary artery disease; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol; NCEP = National Cholesterol Education Program.

<sup>a</sup>Data are presented as number of patients (%) unless otherwise indicated. Includes all patients with more than one visit.

<sup>b</sup>Patients who continued for the full duration of the project.

<sup>c</sup>Patients who did not complete the entire project.

**Figure 3. Persistence, Compliance, and Treatment to NCEP Goal for 397 Patients in Group 1**



ImPACT = Improve Persistence And Compliance with Therapy; NCEP = National Cholesterol Education Program.

## Limitations

The results reported are based on data from 26 sites that continued for the duration of the project. Since the project used an observational, single-cohort design, the results should not be interpreted as proving a cause-and-effect relationship.

## Conclusion

Working collaboratively with patients, physicians, and other health care providers, pharmacists who have ready access to objective clinical data, and the necessary knowledge, skills, and resources, can provide an advanced level of care that results in successful management of dyslipidemia. In this project, mean reductions in both total cholesterol and LDL-C exceeded 30 points for a diverse, multicenter patient population that included both treatment-naïve and previously treated patients who had not achieved goals. Patients enrolled in this project achieved medica-



tion persistence and compliance rates significantly higher than those previously found in the literature from similar ambulatory care settings.

Project ImPACT offers a sound model for pharmacists to use in empowering patients and improving the quality of consumer health outcomes. This approach to health care delivery warrants further investigation and consideration for widespread adoption.

---

Beyond employment by the APhA Foundation of Mr. Bluml, the authors declare no other potential conflicts of interest or financial interests in any product or service mentioned in this article, including grants, employment, gifts, or honoraria.

Project ImPACT: Hyperlipidemia was conducted by the APhA Foundation and was funded through an unrestricted grant from Merck & Co., Inc.

The authors would like to acknowledge Samuel H. Kalman for his support in the development of the project, the interim report, and this report.

## References

- American Heart Association. *2000 Heart and Stroke Statistical Update*. Dallas, Tex: American Heart Association, 1999.
- Castelli WP. Lipoprotein and cardiovascular disease: biological basis and epidemiological studies. Paper presented at: Association for Pharmacoeconomics and Outcomes Research Lipid Conference; November 1997; Orlando, Fla.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Summary of the second report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel II). *JAMA*. 1993;269:3015–23.
- Frick M, Elo MO, Haapa K, et al. Helsinki Heart Study: Primary prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med*. 1987;317:1237–45.
- Lipid Research Clinics Program. The Lipid Research Clinics Coronary Primary Prevention Trial results, I. Reduction in incidence of coronary heart disease. *JAMA*. 1984;251:351–64.
- Sacks FM, Pfeffer MA, Moyer LA, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med*. 1996;335:1001–9.
- Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet*. 1994;344:1383–9.
- Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med*. 1995;333:1301–7.
- Downs JR, Clearfield M, Weis S, et al. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels. Results of AFCAPS/TexCAPS. *JAMA*. 1998;279:1615–22.
- Konzem SL, Gray DR, Kashyap ML. Effect of pharmaceutical care on optimal colestipol treatment in elderly hypercholesterolemic veterans. *Pharmacotherapy*. 1997;17:576–83.
- McKenney JM, Slining JS, Henderson HR, et al. The effect of clinical pharmacy services on patients with essential hypertension. *Circulation*. 1973;48:1104–11.
- Schechtman G, Wolff N, Byrd JC, et al. Physician extenders for cost-effective management of hypercholesterolemia. *J Gen Intern Med*. 1996;11:277–86.
- Health Employer Data Information Set measures [draft]. Reston, Va: National Committee for Quality Assurance; May 1998.
- Clinical Laboratory Improvement Amendments (CLIA '88). P Law No. 100-578, October 31, 1988.
- Hoerger TJ, Vala MV, Bray JW, et al. Treatment patterns and distribution of low-density lipoprotein cholesterol levels in treatment-eligible US adults. *Am J Cardiol*. 1998;82:61–5.
- Avorn J, Monette J, Lacour A, et al. Poor compliance with lipid-lowering medications. *JAMA*. 1998;279:1458–62.
- Simons LA, Levis G, Simons J. Apparent discontinuation rates in patients prescribed lipid-lowering drugs. *Med J Austral*. 1996;164(4):208–11.
- Andrade SE, Walker AM, Gottlieb LK, et al. Discontinuation of antihyperlipidemic drugs—do rates reported in clinical trials reflect rates in primary care settings? *N Engl J Med*. 1995;332:1125–31.
- Eriksson M, Hadell K, Holme I, et al. Compliance with and efficacy of treatment with pravastatin and cholestyramine: a randomized study on lipid-lowering in primary care. *J Intern Med*. 1998;243:373–80.
- Marcelino JJ, Feingold KR. Inadequate treatment with HMG-CoA reductase inhibitors by health care providers. *Am J Med*. 1996;100:605–10.
- Pearson TA, Laurora IM. Treatment success in patient subgroups in the lipid treatment assessment project (L-TAP). *Circulation*. 1998;96: Abstract 361.